

Solutions to In-Class Problems — Week 15, Mon

Problem 1. Prove that the Central Limit Theorem implies the Weak Law of Large Numbers. *Hint:* The only properties of $N(y)$ needed in the proof are that $\lim_{y \rightarrow \infty} N(y) = 1$ and $\lim_{y \rightarrow \infty} N(-y) = 0$.

Solution. Note first that $\mu_{S_n} = n\mu$, $\text{Var}[S_n] = n\sigma^2$, and so $\sigma_{S_n} = \sigma\sqrt{n}$. Now,

$$\begin{aligned} \left| \frac{S_n}{n} - \mu \right| > \epsilon & \text{ iff } |S_n - n\mu| > n\epsilon \\ & \text{ iff } \left| \frac{S_n - n\mu}{\sigma_{S_n}} \right| > \frac{n\epsilon}{\sigma_{S_n}} \\ & \text{ iff } |S_n^*| > \frac{\sqrt{n}\epsilon}{\sigma}. \end{aligned}$$

But for any real number $\beta > 0$,

$$\frac{\sqrt{n}\epsilon}{\sigma} > \beta$$

will hold for all large n . Hence, for any $\beta > 0$ and all large n ,

$$\Pr \left\{ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right\} = \Pr \left\{ |S_n^*| > \frac{\sqrt{n}\epsilon}{\sigma} \right\} \leq \Pr \{ |S_n^*| > \beta \}. \quad (1)$$

So

$$\begin{aligned} \lim_{n \rightarrow \infty} \Pr \left\{ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right\} & \leq \lim_{n \rightarrow \infty} \Pr \{ |S_n^*| > \beta \} && \text{(by (1))} \\ & = \lim_{n \rightarrow \infty} \Pr \{ S_n^* > \beta \} + \Pr \{ S_n^* < -\beta \} \\ & = 1 - N(\beta) + N(-\beta), && \text{(by the Central Limit Thm (8))} \end{aligned}$$

for all real numbers $\beta > 0$. By choosing β large enough, we can ensure that $N(\beta)$ is arbitrarily close to 1 and $N(-\beta)$ is arbitrarily close to 0, so that final term above is arbitrarily close to $1-1+0 = 0$. Hence,

$$\lim_{n \rightarrow \infty} \Pr \left\{ \left| \frac{S_n}{n} - \mu \right| > \epsilon \right\} = 0,$$

which is the Weak Law of Large Numbers. ■

NOTE: We didn't get to the following problem in class.

Problem 2. To clarify the somewhat subtle difference between the Weak and Strong Laws of Large Numbers, we will construct an example of a sequence X_1, X_2, \dots of mutually independent random variables that satisfies the Weak Law of Large Numbers, but not the Strong Law. The distribution of X_i will have to depend on i , because otherwise both laws would be satisfied.¹

In particular, let X_1, X_2, \dots be a sequence of mutually independent random variables such that $X_1 = 0$, and for each integer $i > 1$,

$$\Pr\{X_i = i\} = \frac{1}{2i \log i}, \quad \Pr\{X_i = -i\} = \frac{1}{2i \log i}, \quad \Pr\{X_i = 0\} = 1 - \frac{1}{i \log i}.$$

Note that $\mu = E[X_i] = 0$ for all i .

(a) Show that $\text{Var}[S_n] = \Theta(n^2 / \log n)$. *Hint:* $n / \log n > i / \log i$ for $2 \leq i \leq n$.

Solution.

$$\begin{aligned} \text{Var}[S_n] &= \sum_{i=1}^{\infty} \text{Var}[X_i] && \text{(independent variance additivity)} \\ &= \text{Var}[X_1] + \sum_{i=2}^n E[X_i^2] - E^2[X_i] \\ &= 0 + \sum_{i=2}^n i^2 \left(\frac{1}{i \log i} - 0 \right) \\ &= \sum_{i=2}^n \frac{i}{\log i}. \\ &= \Theta(n^2 / \log n). && \text{(see below)} \quad (2) \end{aligned}$$

To justify (2), note that $x / \log x$ is increasing for $x > e$ since its derivative $(1 / \log x)(1 - 1 / \log x)$ is

¹This problem is adapted from Grinstead & Snell, *Intro. to Probability*, Ch.8, exercise 16, pp314–315, where is credited to David Maslen.

positive. So $n/\log n \geq i/\log i$ for $2 \leq i \leq n$. Therefore,

$$\begin{aligned} \frac{n^2}{\log n} &= \sum_{i=1}^n \frac{n}{\log n} \\ &\geq \sum_{i=2}^n \frac{i}{\log i} \\ &\geq \sum_{i=\lceil n/2 \rceil}^n \frac{i}{\log i} \\ &\geq \sum_{i=\lceil n/2 \rceil}^n \frac{n/2}{\log n} \\ &\geq \frac{n+1}{2} \frac{n/2}{\log n} \\ &\geq \frac{1}{4} \frac{n^2}{\log n}. \end{aligned}$$

■

(b) Show that the sequence X_1, X_2, \dots satisfies the Weak Law of Large Numbers, *i.e.*, prove that for any $\epsilon > 0$

$$\lim_{n \rightarrow \infty} \Pr \left\{ \left| \frac{S_n}{n} \right| \geq \epsilon \right\} = 0.$$

Solution.

$$\begin{aligned} \Pr \left\{ \left| \frac{S_n}{n} \right| \geq \epsilon \right\} &= \Pr \left\{ \left| \frac{S_n}{n} - 0 \right| \geq \epsilon \right\} \\ &\leq \text{Var} \left[\frac{S_n}{n} \right] \frac{1}{\epsilon^2} && \text{(Chebychev Bound)} \\ &= \frac{\text{Var}[S_n]}{n^2} \frac{1}{\epsilon^2} \\ &= \Theta\left(\frac{1}{\epsilon^2 \log n}\right), && \text{(by (2))} \end{aligned}$$

which goes to zero as n goes to ∞ . ■

We now show that the sequence X_1, X_2, \dots does not satisfy the Strong Law of Large Numbers.

(c) (The first Borel-Cantelli lemma.) Let A_1, A_2, \dots be any infinite sequence of mutually independent events such that

$$\sum_{i=1}^{\infty} \Pr \{A_i\} = \infty. \tag{3}$$

Prove that

$$\Pr \{ \text{infinitely many } A_i \text{ occur} \} = 1.$$

Hint: We know that the probability that no A_i with $n \geq i \geq r$ occurs is

$$\leq e^{-E[T_{r,n}]} \quad (4)$$

where $T_{r,n} ::= \sum_{i=r}^n I_{A_i}$ is the number of events A_i with $n \geq i \geq r$ that occur. What happens as $n \rightarrow \infty$?

Solution. Let K_r be the event that no A_i with $i \geq r$ occurs. Also, let $K_{r,n}$ be the event that no A_i with $n \geq i \geq r$ occurs. Finally, let K be the event that only finitely many A_i 's occur. We must prove that $\Pr \{K\} = 0$.

We begin by computing $\lim_{n \rightarrow \infty} e^{-E[T_{r,n}]}$:

$$\begin{aligned} E[T_{r,n}] &= \sum_{i=r}^n E[I_{A_i}] && \text{(linearity of expectation)} \\ &= \sum_{i=r}^n \Pr \{A_i\} && \text{(expectation of indicator variable)} \end{aligned}$$

If we take the the limit as $n \rightarrow \infty$ we have:

$$\begin{aligned} \lim_{n \rightarrow \infty} E[T_{r,n}] &= \lim_{n \rightarrow \infty} \sum_{i=r}^n \Pr \{A_i\} \\ &= \infty && \text{(by (3)).} \end{aligned}$$

Since $e^x \rightarrow 0$ as $x \rightarrow -\infty$, we conclude that:

$$\lim_{n \rightarrow \infty} e^{-E[T_{r,n}]} = 0$$

Note that $K_r \subset K_{r,n}$ for any r, n . Hence $\Pr \{K_r\} \leq \Pr \{K_{r,n}\} \leq e^{-E[T_{r,n}]}$. We have just proved that the right hand side tends to zero as n goes to ∞ . Since the left hand side does not depend on n , and the inequality holds for all n s.t. $n \geq r$, we conclude that $\Pr \{K_r\}$ must be zero.

Now note that $K = \bigcup_r K_r$, so by Boole's law, $\Pr \{K\} \leq \sum_r \Pr \{K_r\}$, and since we've just proved that $\Pr \{K_r\} = 0$ for all r , it follows that $\Pr \{K\} = 0$. Hence the probability that infinitely many A_i 's occur is 1. ■

(d) Show that $\sum_{i=1}^{\infty} \Pr \{|X_i| \geq i\}$ diverges. *Hint:* $\int 1/(x \log x) dx = \log \log x$.

Solution. $\Pr\{|X_i| \geq i\} = 1/(i \log i)$, so

$$\begin{aligned} \sum_{i=1}^n \Pr\{|X_i| \geq i\} &= 0 + \sum_{i=2}^n \frac{1}{i \log i} \\ &\geq \int_2^{n+1} \frac{1}{x \log x} dx \\ &= \log \log(n+1) - \log \log 2, \end{aligned}$$

and this last term approaches infinity as n approaches infinity. ■

(e) Conclude that

$$\Pr\left\{\lim_{n \rightarrow \infty} \frac{S_n}{n} = \mu\right\} = 0. \quad (5)$$

and hence that the Strong Law of Large Numbers *completely* fails for the sequence X_1, X_2, \dots .

Hint:

$$\frac{X_n}{n} = \frac{S_n}{n} - \frac{n-1}{n} \frac{S_{n-1}}{n-1},$$

so if $\lim_{n \rightarrow \infty} S_n/n = 0$, then also $\lim_{n \rightarrow \infty} X_n/n = 0$.

Solution. By parts (c) and (d), the probability that $|X_i| \geq i$ for infinitely many i is 1. But if $|X_i| \geq i$ for infinitely many i , then by definition of the limit, $\lim_{n \rightarrow \infty} X_n/n \neq 0$. Hence,

$$\Pr\left\{\lim_{n \rightarrow \infty} X_n/n \neq 0\right\} = 1,$$

which means

$$\Pr\left\{\lim_{n \rightarrow \infty} X_n/n = 0\right\} = 0, \quad (6)$$

But the hint implies that

$$\Pr\left\{\lim_{n \rightarrow \infty} \frac{S_n}{n} = 0\right\} \leq \Pr\left\{\lim_{n \rightarrow \infty} \frac{X_n}{n} = 0\right\}. \quad (7)$$

Now (6) and (7) immediately imply (5). ■

A Appendix

The *probability density function (pdf)* for a random variable, R , is the function $f_R : \text{range}(R) \rightarrow [0, 1]$ defined by:

$$f_R(x) ::= \Pr\{R = x\}.$$

Random variables R_1, R_2, \dots are *mutually independent* iff

$$\Pr \left\{ \bigcap_i [R_i = x_i] \right\} = \prod_i \Pr \{R_i = x_i\},$$

for all $x_1, x_2, \dots \in \mathbb{R}$. They are *k-wise independent* iff $\{R_i \mid i \in J\}$ are mutually independent for all subsets $J \subset \mathbb{N}$ with $|J| = k$.

Theorem (Weak Law of Large Numbers). Let $S_n ::= \sum_{i=1}^n X_i$, where X_1, \dots, X_n, \dots are pairwise independent variables with the same expectation, μ , and standard deviation, σ . For any $\epsilon > 0$,

$$\lim_{n \rightarrow \infty} \Pr \left\{ \left| \frac{S_n}{n} - \mu \right| \geq \epsilon \right\} = 0.$$

Theorem (The Strong Law of Large Numbers). Let $S_n ::= \sum_{i=1}^n X_i$ where X_1, \dots, X_i, \dots are mutually independent, identically distributed random variables with finite expectation, μ . Then

$$\Pr \left\{ \lim_{n \rightarrow \infty} \frac{S_n}{n} = \mu \right\} = 1.$$

Definition. For any random variable, R , with finite mean, μ_R , and deviation, σ_R , let R^* be the random variable

$$R^* ::= \frac{R - \mu_R}{\sigma_R}.$$

R^* is called the “normalized” version of R .

Definition. The *normal density function* is the function

$$\eta(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2},$$

and the *normal distribution function* is its integral

$$N(y) = \int_{-\infty}^y \eta(x) dx = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^y e^{-x^2/2} dx.$$

The function $\eta(x)$ defines the standard *Bell curve*, centered about the origin with height $1/\sqrt{2\pi}$ and about two-thirds of its area within unit distance of the origin. The normal distribution function $N(y)$ approaches 0 as $y \rightarrow -\infty$. As y approaches zero from below, $N(y)$ grows rapidly towards $1/2$. Then as y continues to increase beyond zero, $N(y)$ rapidly approaches 1.

Theorem (Central Limit Theorem). Let $S_n = \sum_{i=1}^n X_i$ where X_1, \dots, X_i, \dots are mutually independent variables with the same mean, μ , and deviation, σ , and let S_n^* be the normalized version of S_n . Then

$$\lim_{n \rightarrow \infty} \Pr \{S_n^* \leq \beta\} = N(\beta) \tag{8}$$

for any real number β .