

6	9	13	7
12	10	5	
3	4	14	
15	8	11	2

Mathematics for Computer Science

MIT 6.042J/18.062J

Induction II

Copyright © Albert R. Meyer, 2002.

L2-2.1

6	9	13	7
12	10	5	
3	4	14	
15	8	11	2

The MIT Stata Center



<http://web.mit.edu/buildings/statacenter>

Copyright © Albert R. Meyer, 2002.

L2-2.2

6	9	13	7
12	10	5	
3	4	14	
15	8	11	2

The Stata Center Today




Copyright © Albert R. Meyer, 2002.

L2-2.3

6	9	13	7
12	10	5	
3	4	14	
15	8	11	2

The Stata Center Plaza




Copyright © Albert R. Meyer, 2002.

L2-2.4

6	9	13	7
12	10	5	
3	4	14	
15	8	11	2

The Gehry/Gates Plaza

Goal: tile the squares, except one in the middle for Bill.



Copyright © Albert R. Meyer, 2002.

L2-2.5


6	9	13	7
12	10	5	
3	4	14	
15	8	11	2

The Gehry/Gates Plaza

Gehry specifies L-shaped tiles covering three squares:



For example, for 8 x 8 plaza might tile for Bill this way:



Copyright © Albert R. Meyer, 2002.

L2-2.6




The Gehry/Gates Plaza

Theorem: For any $2^n \times 2^n$ plaza, we can make Bill happy.


Proof: (by induction on n)
 $P(n) ::=$ can tile $2^n \times 2^n$ with Bill in middle.

Base case: ($n=0$)




The Gehry/Gates Plaza

Induction step: assume can tile $2^n \times 2^n$,
prove can handle $2^{n+1} \times 2^{n+1}$.




The Gehry/Gates Plaza

Method: Divide into subsquares
Relocate the subsquares as indicated.




The Gehry/Gates Plaza

Method: after relocation have:
Now rotate the squares as indicated.



The Gehry/Gates Plaza

Method: after rotation have:
Now put tile in center



The Gehry/Gates Plaza


Did I fool you?



The Gehry/Gates Plaza

Bug:

Division into subsquares may



The Gehry/Gates Plaza

The fix:

Prove that we can always find a tiling with Bill **in the corner**.




The Gehry/Gates Plaza

Induction step:


Assume we can get Bill **in corner** of $2^n \times 2^n$.

Prove we can get Bill in corner of $2^{n+1} \times 2^{n+1}$.




The Gehry/Gates Plaza

Method: Rotate the squares as indicated.




The Gehry/Gates Plaza

Method: Rotate the squares as indicated.
after rotation have:



The Gehry/Gates Plaza


Method: Now group the squares together,
and fill the center with a tile.



6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

The Gehry/Gates Plaza

Note: Once have Bill in corner, can get Bill in middle:




Copyright © Albert R. Meyer, 2002.

L2-2.19

6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

The Gehry/Gates Plaza

Method:
Rotate the squares as indicated.




Copyright © Albert R. Meyer, 2002.

L2-2.20

6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

The Gehry/Gates Plaza

Method: after rotation have:




Copyright © Albert R. Meyer, 2002.

L2-2.21

6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

The Gehry/Gates Plaza

Method: Now group the 4 squares together, and insert a tile.



Copyright © Albert R. Meyer, 2002.

L2-2.22

6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

Ingenious Induction Hypotheses

Note 1: To prove
“Bill in middle,”
we *proved something else*: “Bill in corner.”

Copyright © Albert R. Meyer, 2002.

L2-2.23

6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

Inductive (Recursive) Procedures

Note 2: The induction proof of
“Bill in corner” implicitly defines
a *recursive procedure* for
constructing a $2^{n+1} \times 2^{n+1}$ corner
tiling from a $2^n \times 2^n$ tiling.

Copyright © Albert R. Meyer, 2002.

L2-2.24

6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

Ingenious Induction Hypotheses

Note 3: Other times it helps to choose a *stronger hypothesis* than the desired result. Result at $n+1$ becomes harder to prove -- but we have a stronger hypothesis at n to prove it with!

6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

Problems

Class Problem 1

6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

A False Proof

Theorem: All horses are the same color.

Proof: (by induction on n)

Induction hypothesis:

$P(n) ::=$ any set of n horses have the same color

Base case ($n=0$):

No horses so *vacuously* true!




6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

A False Proof

(Inductive case)

Assume any n horses have the same color.

Prove that any $n+1$ horses have the same color.




6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

A False Proof

(Inductive case)

Assume any n horses have the same color.

Prove that any $n+1$ horses have the same color.




6	9	13	7
12	10	5	
3	4	14	11
15	8	16	2

A False Proof

(Inductive case)

Assume any n horses have the same color.

Prove that any $n+1$ horses have the same color.



6	9	13	7
12	10	8	
3	4	14	5
15	11	2	1

A False Proof

What is wrong? $n=1$

Proof that $P(n) \rightarrow P(n+1)$
is **false** if $n=1$, because the two
horse groups *do not overlap*.

Second set of $n=1$ horses



First set of $n=1$ horses

Copyright © Albert R. Meyer, 2002.

L2-2.31

6	9	13	7
12	10	8	
3	4	14	5
15	11	2	1

A False Proof

Proof that $P(n) \rightarrow P(n+1)$
is **false** if $n=1$, because the two
horse groups *do not overlap*.

(But proof works for all $n \neq 1$)

Copyright © Albert R. Meyer, 2002.

L2-2.32

6	9	13	7
12	10	8	
3	4	14	5
15	11	2	1

Problems

Class Problems 2 & 3

Copyright © Albert R. Meyer, 2002.

L2-2.33